

Exact eigenvalue spectrum of a class of fractal scale-free networks

ZHONGZHI ZHANG^{1,2 (a)}, ZHENGYI HU^{1,2}, YIBIN SHENG³ and GUANRONG CHEN⁴

¹ School of Computer Science, Fudan University, Shanghai 200433, China

² Shanghai Key Lab of Intelligent Information Processing, Fudan University, Shanghai 200433, China

³ School of Mathematical Sciences, Fudan University, Shanghai 200433, China

⁴ Department of Electronic Engineering, City University of Hong Kong, Hong Kong SAR, China

PACS 05.40.Fb – Random walks and Levy flights
 PACS 89.75.Hc – Networks and genealogical trees
 PACS 02.10.Yn – Matrix theory

Abstract – The eigenvalue spectrum of the transition matrix of a network encodes important information about its structural and dynamical properties. We study the transition matrix of a family of fractal scale-free networks and analytically determine all the eigenvalues and their degeneracies. We then use these eigenvalues to evaluate the closed-form solution to the eigentime for random walks on the networks under consideration. Through the connection between the spectrum of transition matrix and the number of spanning trees, we corroborate the obtained eigenvalues and their multiplicities.

Introduction. – Complex networks have emerged as a universal tool for studying complex systems in many different fields, such as physics, chemistry, biology, and computer science [1–3]. A fundamental issue in network science is to understand diverse dynamical processes defined on different networks [4–6]. A paradigmatic dynamical process that has attracted increasing research interest is random walks [7–17], which have found a broad range of applications in various areas of science and engineering [18–21].

As is well known, random walks on a network can be described by a transition matrix and various interesting quantities about random walks are closely related to the spectrum of the transition matrix [17, 22–24]. For example, the mean first-passage time (MFPT) from one node to another on a network can be expressed in terms of the eigenvalues and orthonormalized eigenvectors of its transition matrix [17, 22, 23]; the sum of reciprocals of one minus each eigenvalue, other than the eigenvalue 1, determines the eigentime identity [23], sometime referred to as Kemeny’s constant [24]; the second largest eigenvalue, together with the smallest eigenvalue, determines the mix time of random walks on the network [22, 23]. Thus, it is of theoretical and practical significance to calculate and analyze the spectra of the transition matrices of networks. However, little attention has been paid to this relevant is-

sue, the main reason for which is that the computational complexity for computing the eigenvalue spectrum of a general network is very high [25].

In this paper, we present a theoretical study of the transition matrix for unbiased random walks on a family of iterative networks [26–28], which have remarkable scale-free [29] and fractal [30] phenomena that constitute our fundamental understanding of real-life systems. Making use of the real-space decimation technique [31, 32], we will derive an exact recursion expression for the eigenvalues at every two consecutive iterations, which is obtained based on the structure of the studied networks. We then proceed to provide the complete spectrum of eigenvalues and their corresponding degeneracies. Using these eigenvalues, we further derive an explicit formula for the eigentime identity. Finally, we prove the validity of the computation for eigenvalues based on the relationship between the number of spanning trees and the eigenvalues of the transition matrix of a network.

Model and properties. – The considered fractal scale-free networks are constructed in an iterative way [26–28]. Let F_n ($n \geq 0$) denote the networks after n iterations. For $n = 0$, F_0 consists of an edge connecting two nodes. For $n \geq 1$, F_n is obtained from F_{n-1} by performing the following operations on every existing edge in F_{n-1} as shown in Fig. 1: replace the edge by a path of 2 links long, with both endpoints of the path being the iden-

^(a) zhangzz@fudan.edu.cn




Fig. 1: (Color online) Construction of the networks. One can obtain the next generation of the network family through replacing each edge of the present generation by the clusters on the right-hand side of the arrow.

tical endpoints of the original edge, then for each endpoint of the path create m (a positive integer) new nodes and attach them to the endpoint. Figure 2 illustrates the first several iterative construction steps for a particular case of $m = 1$, whose dynamics of random walks were studied in [33].

The deterministic construction of the networks allows for treating exactly their relevant properties. At each generation n_i ($n_i \geq 1$), the number of newly introduced nodes is $V_{n_i} = (2m + 1)(2m + 2)^{n_i - 1}$. Then, the network size (number of nodes) N_n of F_n is given by

$$N_n = \sum_{n_i=0}^n V_{n_i} = (2m + 2)^n + 1. \quad (1)$$

And the total number of edges is

$$E_n = V_n - 1 = (2m + 2)^n. \quad (2)$$

Let $d_i(n)$ be the degree of node i in F_n that entered the networks at generation n_i ($n_i \geq 0$). Then $d_i(n + 1) = (m + 1)d_i(n)$, that is, after each new iteration the degree of node i increases by m times.

These resultant networks display the following interesting structural features. They are in power law with the degree distribution exponent being $\gamma = 1 + \ln(2m + 2)/\ln(m + 1)$ [27, 34, 35]. In addition, after the evolution of one generation, the number of nodes increases by a factor $f_N = 2m + 2$, see Eq. (1); the diameter grows by a factor of $f_D = 2$ [27], and the MFPT between two previously existing nodes increases by a factor of $f_M = 4m + 4$ [27, 35]. Thus, the networks are fractal with a fractal dimension $f_B = \ln f_N / \ln f_D = \ln(2m + 2) / \ln 2$ [11, 26, 34]; their random-walk dimension is $f_w = \ln f_M / \ln f_D = \ln(4m + 4) / \ln 2$, and their spectral dimension is $f_s = 2f_B / f_w = 2 \ln(2m + 2) / \ln(4m + 4)$. Finally, they are “large-world” with their diameter and average distance growing in a power of the network size [27, 34, 35].

Spectrum of transition matrix. – After introducing the model and some properties of the networks, in this section we study the eigenvalue spectrum of the transition matrix for the network family, as well as their degeneracies.

Eigenvalue spectrum. As is well known, the structure of F_n is encoded in its adjacency matrix A_n , whose entries $A_n(i, j)$ are defined by $A_n(i, j) = 1$ if nodes i and




Fig. 2: (Color online) Growth process for a special case of network corresponding to $m = 1$.

j are adjacent in F_n , or $A_n(i, j) = 0$ otherwise. Then, the transition matrix of F_n , denoted by T_n , is defined as $T_n = D_n^{-1} A_n$, where D_n is the diagonal degree matrix of F_n with its i th diagonal entry being $d_i(n)$. Thus, the element of T_n is $T_n(i, j) = A_n(i, j) / d_i(n)$, which represents the jumping probability of the discrete-time unbiased random walks [7] for a particle going from node i to node j .

We now consider the eigenvalue spectrum of T_n . Since T_n is asymmetric, we introduce the following matrix:

$$P_n = D_n^{-\frac{1}{2}} A_n D_n^{-\frac{1}{2}} = D_n^{\frac{1}{2}} T_n D_n^{-\frac{1}{2}}. \quad (3)$$

Obviously, P_n is real and symmetric and has the same set of eigenvalues as T_n . It is easy to verify that the entry of P_n is $P_n(i, j) = \frac{A_n(i, j)}{\sqrt{d_i(n)} \sqrt{d_j(n)}}$. Next, we apply the decimation method [31, 32] to determine the eigenvalues and their multiplicities of P_n . The decimation approach is universal and has been used to compute the Laplacian spectra of Vicsek fractals [36–38] and their extensions [39].

We now address the eigenvalue problem for matrix P_{n+1} . Let α represent the set of nodes belonging to F_n , and β the set of nodes generated at $(n + 1)$ th iteration. By definition, P_{n+1} has a block form

$$P_{n+1} = \begin{bmatrix} P_{\alpha, \alpha} & P_{\alpha, \beta} \\ P_{\beta, \alpha} & P_{\beta, \beta} \end{bmatrix} = \begin{bmatrix} 0 & P_{\alpha, \beta} \\ P_{\beta, \alpha} & 0 \end{bmatrix}, \quad (4)$$

where we have used the facts that $P_{\alpha, \alpha}$ is the zero matrix of order $N_n \times N_n$ since there are no transitions between any pair of original nodes, and that $P_{\beta, \beta}$ is the zero matrix of order $(N_{n+1} - N_n) \times (N_{n+1} - N_n)$.

Suppose $\lambda_i(n + 1)$ is an eigenvalue of P_{n+1} , and $u = (u_\alpha, u_\beta)^\top$ is its associated eigenvector, where the superscript $\top$ represents transpose and u_α and u_β correspond to nodes in α and β , respectively. Then, one can write the eigenvalue equation for matrix P_{n+1} in the following block form:

$$\begin{bmatrix} 0 & P_{\alpha, \beta} \\ P_{\beta, \alpha} & 0 \end{bmatrix} \begin{bmatrix} u_\alpha \\ u_\beta \end{bmatrix} = \lambda_i(n + 1) \begin{bmatrix} u_\alpha \\ u_\beta \end{bmatrix}. \quad (5)$$

Equation (5) can be recast into two equations:

$$P_{\alpha, \beta} u_\beta = \lambda_i(n + 1) u_\alpha, \quad (6)$$

$$P_{\beta,\alpha}u_\alpha = \lambda_i(n+1)u_\beta. \quad (7)$$

Equation (7) yields

$$u_\beta = \frac{1}{\lambda_i(n+1)}P_{\beta,\alpha}u_\alpha, \quad (8)$$

provided that $\lambda_i(n+1) \neq 0$. Plugging Eq. (8) into Eq. (6) gives

$$\frac{1}{\lambda_i(n+1)}P_{\alpha,\beta}P_{\beta,\alpha}u_\alpha = \lambda_i(n+1)u_\alpha. \quad (9)$$

Thus, we reduce the problem of determining the eigenvalues for matrix P_{n+1} of order $N_{n+1} \times N_{n+1}$ to calculating the eigenvalues of matrix $P_{\alpha,\beta}P_{\beta,\alpha}$ with a smaller order of $N_n \times N_n$.

In the Appendix, we prove that

$$P_{\alpha,\beta}P_{\beta,\alpha} = \frac{2m+1}{2m+2}I_n + \frac{1}{2m+2}P_n, \quad (10)$$

where I_n is the identity matrix of the same order as P_n . Thus, we have related $P_{\alpha,\beta}P_{\beta,\alpha}$ to P_n , which enables us to express the eigenvalues of matrix P_{n+1} in terms of those of P_n .

Substituting Eq. (10) into Eq. (9), we obtain

$$\frac{1}{\lambda_i(n+1)}\left(\frac{2m+1}{2m+2}I_n + \frac{1}{2m+2}P_n\right)u_\alpha = \lambda_i(n+1)u_\alpha, \quad (11)$$

that is,

$$P_n u_\alpha = \{(2m+2)[\lambda_i(n+1)]^2 - (2m+1)\}u_\alpha. \quad (12)$$

Hence, if $\lambda_i(n)$ is the eigenvalue of P_n corresponding to the eigenvector u_α , Eq. (12) implies

$$\lambda_i(n) = (2m+2)[\lambda_i(n+1)]^2 - (2m+1). \quad (13)$$

Solving the quadratic equation in the variable $\lambda_i(n+1)$ given by Eq. (13) yields

$$\lambda_{i,1}(n+1) = \sqrt{\frac{2m+1+\lambda_n}{2m+2}}, \quad \lambda_{i,2}(n+1) = -\lambda_{i,1}(n+1). \quad (14)$$

Equation (14) relates $\lambda_i(n+1)$ to $\lambda_i(n)$, with each $\lambda_i(n)$ giving rise to two eigenvalues of P_{n+1} .

Degeneracies of eigenvalues. To determine the multiplicities of the eigenvalues, we first calculate numerically the eigenvalues for those networks of small sizes. For F_0 , the eigenvalues are 1 and -1 ; while for F_1 , its eigenvalues are 1, -1 , 0, $\frac{\sqrt{m+m^2}}{m+1}$, and $\frac{-\sqrt{m+m^2}}{m+1}$. For $n \geq 2$, we find that the eigenvalue spectrum displays the following properties: (i) From Eq. (14), eigenvalue 1 gives rise to two eigenvalues 1 and -1 with single degeneracy. (ii) All eigenvalues of a given generation n_i always exist at its subsequent generation $n_i + 1$, and all new eigenvalues at generation $n_i + 1$ are just those generated via Eq. (14) by substituting $\lambda_i(n)$ with $\lambda_i(n_i)$ that are newly added to generation n_i ; moreover each new eigenvalue keeps the

degeneracy of its father. (iii) Except for 0, all other eigenvalues are generated from -1 and 0, and all the offspring eigenvalues of -1 are nondegenerate. Thus, all that is left is to determine the degeneracy of 0 and the multiplicities of its offsprings, based on property (ii).

Let $D_n^{\text{mul}}(\lambda)$ denote the multiplicity of eigenvalue λ of matrix P_n . We now find the number of eigenvalue 0 of P_n . Let $r(M)$ be the rank of matrix M . Then, the degeneracy of eigenvalue 0 of P_{n+1} is

$$D_{n+1}^{\text{mul}}(\lambda = 0) = N_{n+1} - r(P_{n+1}). \quad (15)$$

In order to determine $D_{n+1}^{\text{mul}}(\lambda = 0)$, we can alternatively compute $r(P_{n+1})$. Obviously, $r(P_{n+1}) = r(P_{\alpha,\beta}) + r(P_{\beta,\alpha}) = 2r(P_{\beta,\alpha})$, where $r(P_{\alpha,\beta}) = r(P_{\beta,\alpha})$ is used.

We next determine $r(P_{\beta,\alpha})$. First, we show that $P_{\beta,\alpha}$ is a full column rank matrix. Let

$$v = (v_1, v_2, \dots, v_{N_{n+1}-N_n})^\top = \sum_{\substack{i=1 \\ i \in \alpha}}^{N_n} k_i M_i, \quad (16)$$

where M_i is the column vector of $P_{\beta,\alpha}$ as defined in Eq. (36). Let $M_i = (M_{1,i}, M_{2,i}, \dots, M_{N_{n+1}-N_n,i})^\top$. Suppose that $v = 0$. Then, we can prove that for an arbitrary k_i , $k_i = 0$ always holds. By construction, for any old node $i \in \alpha$, there exists a new leaf node $l \in \beta$ attached to i . Then, for $v_l = k_1 M_{1,l} + k_2 M_{2,l} + \dots + k_{N_{n+1}-N_n,l} M_{N_{n+1}-N_n,l}$, only $M_{i,l} \neq 0$ but all $M_{x,l} = 0$ for $x \neq i$. From $v_l = 0$, we have $k_i = 0$. Therefore, $r(P_{\beta,\alpha}) = N_n = (2m+2)^n + 1$.

Combining the above-obtained results, we obtain the degeneracy of eigenvalue 0 of P_n as

$$D_n^{\text{mul}}(\lambda = 0) = \begin{cases} 0, & n = 0, \\ 2m(2m+2)^{n-1} - 1, & n \geq 1. \end{cases} \quad (17)$$

Since every eigenvalue in P_n keeps the degeneracy of its father, the multiplicity of each first-generation descendants of eigenvalue 0 is $2m(2m+2)^{n-2} - 1$, the multiplicity of each second-generation descendants of eigenvalue 0 is $2m(2m+2)^{n-3} - 1$, and so on. Thus, the total number of eigenvalue 0 and all of its descendants in P_n ($n \geq 1$) is

$$\begin{aligned} N_n^{\text{seed}}(0) &= \sum_{i=1}^n [2m(2m+2)^{i-1} - 1] 2^{n-i} \\ &= (2m+2)^n - 2^{n+1} + 1. \end{aligned} \quad (18)$$

Similarly, the total number of eigenvalue -1 and its descendants in P_n ($n \geq 0$) is

$$N_n^{\text{seed}}(-1) = \sum_{i=0}^n 2^i = 2^{n+1} - 1. \quad (19)$$

Summing up the number of eigenvalues found above, we have

$$N_n^{\text{seed}}(0) + N_n^{\text{seed}}(-1) + 1 = (2m+2)^n + 1 = N_n, \quad (20)$$

which means that we have determined all the eigenvalues of P_n .

Application of eigenvalue spectrum. — We next show how to use the obtained eigenvalues and their multiplicities to determine some related quantities for the fractal scale-free network family F_n , including the eigentime identity of random walks and the number of spanning trees. We note that the number of spanning trees for every generation is 1, so our aim for evaluating spanning trees is to verify that our computation of the eigenvalues of the transition matrix is right.

Eigentime identity. Eigentime identity of random walks on a network is a global characteristic of the network, which reflects the structure of the whole network.

Let $H_{ij}(n)$ denote the mean-first passage time from node i to node j in F_n , which is the expected time for a walker starting from node i to reach node j for the first time. Let $\pi = (\pi_1, \pi_2, \dots, \pi_N)^\top$ represent the stationary distribution for random walks on F_n [22, 23]. It is easy to derive $\pi_i = d_i(n)/(2E_n)$, satisfying $\sum_{i=1}^N \pi_i = 1$ and $\pi^\top T_n = \pi^\top$. Then, the eigentime identity, denoted by H_n , for random walks on F_n , is the expected time from a node i to another node j , selected randomly from all nodes accordingly to the stationary distribution. That is,

$$H_n = \sum_{j=1}^{N_n} \pi_j H_{ij}(n) = \sum_{i=2}^{N_n} \frac{1}{1 - \lambda_i(n)}, \quad (21)$$

where we have assumed that $\lambda_1(n) = 1$.

Note that for any i , $\sigma_i(n) = 1 - \lambda_i(n)$ is an eigenvalue of the normalized Laplacian matrix of F_n , defined as $L_n = I_n - P_n$ [40–42]. The normalized Laplacian matrix has found numerous applications [43–45]. By definition,

$$H_n = \sum_{i=2}^{N_n} \frac{1}{\sigma_i(n)}. \quad (22)$$

The goal next is to explicitly evaluate this sum.

Let Ω_n be the set of all the $N_n - 1$ nonzero eigenvalues of matrix L_n , $\Omega_n = \{\sigma_2(n), \sigma_3(n), \dots, \sigma_{N_n}(n)\}$, in which the distinctness of the elements has been ignored. It is clear that Ω_n ($n \geq 1$) includes 1, 2, and other eigenvalues generated by them. Then, Ω_n can be classified into three subsets represented by $\Omega_n^{(1)}$, $\Omega_n^{(2)}$ and $\Omega_n^{(3)}$, respectively. That is, $\Omega_n = \Omega_n^{(1)} \cup \Omega_n^{(2)} \cup \Omega_n^{(3)}$, where $\Omega_n^{(1)}$ consists of eigenvalue 1 with multiplicity $2m(2m+2)^{n-1} - 1$, $\Omega_n^{(2)}$ contains only eigenvalue 2 with a single degeneracy, and $\Omega_n^{(3)}$ includes those eigenvalues generated by 1 and 2. Obviously, $\sum_{i \in \Omega_n^{(1)}} \frac{1}{\sigma_i(n)} = 2m(2m+2)^{n-1} - 1$ and $\sum_{i \in \Omega_n^{(2)}} \frac{1}{\sigma_i(n)} = \frac{1}{2}$.

Each eigenvalue $\sigma_i(n-1)$ in $\Omega_{n-1}^{(3)}$ generates two eigenvalues, $\sigma_{i,1}(n)$ and $\sigma_{i,2}(n)$, belonging to $\Omega_n^{(3)}$, via the following equation:

$$[\sigma_i(n)]^2 - 2\sigma_i(n) + \frac{\sigma_i(n-1)}{2(m+1)} = 0, \quad (23)$$

which is easily obtained from Eq. (13). According to Vieta's formulas, we have $\sigma_{i,1}(n) + \sigma_{i,2}(n) = 2$ and $\sigma_{i,1}(n) \times \sigma_{i,2}(n) = \frac{\sigma_i(n-1)}{2(m+1)}$. Then

$$\frac{1}{\sigma_{i,1}(n)} + \frac{1}{\sigma_{i,2}(n)} = \frac{4(m+1)}{\sigma_i(n-1)}, \quad (24)$$

which implies that

$$\sum_{\sigma_i(n) \in \Omega_n^{(3)}} \frac{1}{\sigma_i(n)} = 4(m+1) \sum_{\sigma_i(n-1) \in \Omega_{n-1}^{(3)}} \frac{1}{\sigma_i(n-1)}. \quad (25)$$

Thus, we have

$$H_n = 4(m+1)H_{n-1} + [2m(2m+2)^{n-1} - 1] + \frac{1}{2}. \quad (26)$$

Considering the initial condition $H_1 = \frac{8m+3}{2}$, Eq. (26) can be solved by induction to yield

$$H_n = \frac{(6m^2 + 6m + 1)(4m + 4)^n}{(m+1)(4m+3)} - 2m(2m+2)^{n-1} + \frac{1}{8m+6}, \quad (27)$$

which can be expressed in terms of network size N_n as

$$H_n = \frac{6m^2 + 6m + 1}{(m+1)(4m+3)} (N_n - 1)^{1 + \ln 2 / \ln(2m+2)} - \frac{m}{m+1} (N_n - 1) + \frac{1}{8m+6}. \quad (28)$$

Thus, for large networks, i.e., $N_g \rightarrow \infty$,

$$H_n \sim (N_n)^{1 + \ln 2 / \ln(2m+2)} = (N_n)^{1 + 1/f_B}, \quad (29)$$

increasing as a power-law function of the network size N_n with the power exponent greater than 1. Note that the power-law exponent in Eq. (29) is consistent with the general scaling for the average global MFPT (i.e., eigentime identity) previously obtained in [13], but is comparable to the diverse behaviors of MFPT observed in real-life systems, the reason for which was explained in [46].

Spanning trees. An important task in the study of networks is to determine the number of spanning trees on different networks [47–51]. A spanning tree of a connected graph is defined as a maximal set of edges of the graph that contains no cycle, or as a minimal set of edges that connect all nodes. It has been shown [40, 42] that for a connected network G with N nodes, the number of its spanning trees, $N_{\text{st}}(G)$, is related to the $N - 1$ nonzero eigenvalues of its normalized Laplacian matrix. According to [40, 42], the number of spanning trees $N_{\text{st}}(F_n)$ for F_n is determined by

$$N_{\text{st}}(F_n) = \prod_{i=1}^{N_n} d_i(n) \prod_{i=2}^{N_n} \sigma_i(n) \Big/ \sum_{i=1}^{N_n} d_i(n). \quad (30)$$

We now determine the terms in the sum and in the products of Eq. (30). It is evident that

$$\sum_{i=1}^{N_n} d_i(n) = 2E_n = (2m+2)^n. \quad (31)$$

Let Δ_n and Λ_n denote $\prod_{i=1}^{N_n} d_i(n)$ and $\prod_{i=2}^{N_n} \sigma_i(n)$, respectively. According to the above-obtained results, it is easy to show that they obey the following recursive relations:

$$\Delta_n = \Delta_{n-1} \times (m+1)^{N_{n-1}} \times 2^{E_{n-1}} \quad (32)$$

and

$$\Lambda_n = 2 \left(\frac{1}{2m+2} \right)^{N_{n-1}-1} \Lambda_{n-1}. \quad (33)$$

Using $\Delta_1 = 1$ and $\Lambda_1 = 2$, we can solve Eqs. (32) and (33) to obtain

$$\Delta_n = (m+1)^n (2m+2)^{\frac{(2m+2)^n - 1}{2m+1}} \quad (34)$$

and

$$\Lambda_n = 2^{n+1} \left(\frac{1}{2m+2} \right)^{\frac{(2m+2)^n - 1}{2m+1}}. \quad (35)$$

Inserting Eqs. (31), (34) and (35) into Eq. (30) gives $N_{\text{st}}(F_n) = 1$, which proves that the technique and process of our computation on the eigenvalues and their degeneracies for the transition matrix of the network family F_n are indeed correct.

Conclusions. – Rich information about the structure and random-walk dynamics of a network can be extracted from the spectrum of the transition matrix of the network. In this paper, we have studied the transition matrix of a class of networks with scale-free and fractal behaviors that are observed in various real-world networked systems. The iterative construction of the networks allows a detailed analysis of the eigenvalues and their multiplicities. Using the renormalization approach, we have derived an explicit recursive relationship between the eigenvalues of the network family at two successive iterations. For each eigenvalue, we have also determined its degeneracy. Based on the obtained recursion relation of eigenvalues, we have further determined the eigentime identity for random walks, an important quantity rooted in the inherent topology of a network. Moreover, we have tested our computation and results for the spectrum of the transition matrix via evaluating the number of spanning trees in the network family under consideration, using the established relation between the two quantities. Our work opens up the possibilities to characterize the spectrum for the transition matrices of other deterministic networks.

It deserves to be mentioned that by definition eigentime identity is the expected time from a node i to another node j that is chosen randomly from all nodes according to the stationary distribution, which means that the target is immobile. Then, interesting questions arise: What happens if the target is not immobile? Would it be possible in this case to use the so-called Pascal Principle [13, 52, 53] to obtain a rigorous upper bound? Although the questions go beyond the scope of the present paper, they are worth studying in the future. Furthermore, future work should also include how to apply the theory to real-world networks, a relation between whose topology and function has been demonstrated [54].

We would like to thank Bin Wu for assistance. This work was supported by the National Natural Science Foundation of China under Grant No. 61074119 and the Hong Kong Research Grants Council under the GRF Grant CityU 1114/11E.

Proof of equation (10). – In order to prove the equivalent relation $P_{\alpha,\beta} P_{\beta,\alpha} = \frac{2m+1}{2m+2} I_n + \frac{1}{2m+2} P_n$, it suffices to show that the entries of the former are equal to their counterparts of the latter. For matrix $\frac{2m+1}{2m+2} I_n + \frac{1}{2m+2} P_n$, denoted by Q_n , obviously its entries are: $Q_n(i, i) = \frac{2m+1}{2m+2}$ for $i \neq j$ and $Q_n(i, j) = \frac{1}{2m+2} P_n(i, j)$ otherwise.

We proceed to determine the entries of $P_{\alpha,\beta} P_{\beta,\alpha}$, denoted by R_n . Note that $P_{\alpha,\beta}$ can be written as

$$P_{\alpha,\beta} = \begin{pmatrix} M_1^\top \\ M_2^\top \\ \vdots \\ M_{N_n}^\top \end{pmatrix}, \quad (36)$$

where each M_i is a column vector with order $N_{n+1} - N_n$. Since $P_{\alpha,\beta} = P_{\beta,\alpha}^\top$, we have $P_{\beta,\alpha} = (M_1 M_2 \dots M_{N_n})$. Then, the entry $R_n(i, j) = M_i^\top M_j$ of R_n can be determined in the following way.

If $i = j$, the diagonal element of R_n is

$$\begin{aligned} R_n(i, i) &= M_i^\top M_i \\ &= \sum_{k \in \beta} P_{n+1}(i, k) P_{n+1}(k, i) = \sum_{k \in \beta} \frac{A_{n+1}(i, k)}{d_i(n+1) d_k(n+1)} \\ &= \frac{1}{d_i(n+1)} \left(\sum_{\substack{k \in \beta, i \sim k \\ d_k(n+1)=2}} \frac{1}{2} + \sum_{\substack{k \in \beta, i \sim k \\ d_k(n+1)=1}} 1 \right) \\ &= \frac{1}{d_i(n+1)} \left[d_i(n) \times \frac{1}{2} + m d_i(n) \right] \\ &= \frac{2m+1}{2m+2} = Q_n(i, i), \end{aligned} \quad (37)$$

where the relation $d_i(n+1) = (m+1) d_i(n)$ has been used. In Eq. (37), $i \sim k$ means that nodes i and k are adjacent in F_{n+1} .

If $i \neq j$, the non-diagonal element of R_n is

$$\begin{aligned} R_n(i, j) &= M_i^\top M_j \\ &= \sum_{k \in \beta} P_{n+1}(i, k) P_{n+1}(k, j) \\ &= \sum_{\substack{A_{n+1}(i,k)=1 \\ A_{n+1}(k,j)=1}} \frac{A_n(i, j)}{d_k(n+1) \sqrt{d_i(n+1) d_j(n+1)}} \\ &= \frac{A_n(i, j)}{(2m+1) \sqrt{d_i(n) d_j(n)}} = Q_n(i, j). \end{aligned} \quad (38)$$

Thus, Eq. (10) is proved.

REFERENCES

- [1] R. Albert and A.-L. Barabási, *Rev. Mod. Phys.* **74**, 47 (2002).
- [2] S. N. Dorogovtsev and J. F. F. Mendes, *Adv. Phys.* **51**, 1079 (2002).
- [3] M. E. J. Newman, *Networks: An Introduction* (Oxford University Press, New York, 2010).
- [4] M. E. J. Newman, *SIAM Rev.* **45**, 167 (2003).
- [5] S. Boccaletti, V. Latora, Y. Moreno, M. Chavez and D.-U. Hwang, *Phys. Rep.* **424**, 175 (2006).
- [6] S. N. Dorogovtsev, A. V. Goltsev and J.F.F. Mendes, *Rev. Mod. Phys.* **80**, 1275 (2008).
- [7] J. D. Noh and H. Rieger, *Phys. Rev. Lett.* **92**, 118701 (2004).
- [8] V. Sood, S. Redner, and D. ben-Avraham, *J. Phys. A* **38**, 109 (2005).
- [9] E. M. Bollt, D. ben-Avraham, *New J. Phys.* **7**, 26 (2005).
- [10] S. Condamin, O. Bénichou, V. Tejedor, R. Voituriez, and J. Klafter, *Nature (London)* **450**, 77 (2007).
- [11] L. K. Gallos, C. Song, S. Havlin, and H. A. Makse, *Proc. Natl. Acad. Sci. USA* **104**, 7746 (2007).
- [12] A. Kittas, S. Carmi, S. Havlin, and P. Argyrakis, *EPL* **84**, 40008 (2008).
- [13] V. Tejedor, O. Bénichou, and R. Voituriez, *Phys. Rev. E* **80**, 065104(R) (2009).
- [14] A. Fronczak and P. Fronczak, *Phys. Rev. E* **80**, 016107 (2009).
- [15] A. P. Roberts and C. P. Haynes, *Phys. Rev. E* **83**, 031113 (2011).
- [16] V. Tejedor, O. Bénichou, and R. Voituriez, *Phys. Rev. E* **83**, 066102 (2011).
- [17] Z. Z. Zhang, J. L. T. Alafate, B. Y. Hou, H. J. Zhang, and G. R. Chen, *Eur. Phys. J. B* **84**, 691 (2011).
- [18] G. H. Weiss, *Aspects and Applications of the Random Walk* (North Holland, Amsterdam, 1994).
- [19] B. H. Hughes, *Random Walks and Random Environments* (Clarendon Press, Oxford, 1995), Vols. 1 and 2.
- [20] S. Redner, *A Guide to First-Passage Processes* (Cambridge University Press, Cambridge, England 2001).
- [21] O. Bénichou, C. Loverdo, M. Moreau, and R. Voituriez, *Rev. Mod. Phys.* **83**, 81 (2011).
- [22] L. Lovász, *Random Walks on Graphs: A Survey*, in *Combinatorics, Paul Erdős is Eighty Vol. 2*, edited by D. Miklós, V. T. Sós, and T. Szőnyi (János Bolyai Mathematical Society, Budapest, 1996), pp. 353-398; <http://research.microsoft.com/users/lovasz/erdos.ps>.
- [23] D. Aldous and J. Fill, *Reversible Markov chains and random walks on graphs*, 1999, <http://www.stat.berkeley.edu/~aldous/RWG/book.html>.
- [24] M. Levene and G. Loizou, *Am. Math. Mon.* **109**, 741 (2002).
- [25] N. L. Biggs, *Algebraic Graph Theory*, 2nd ed. (Cambridge University Press, Cambridge, 1993).
- [26] C. Song, S. Havlin, H. A. Makse, *Nat. Phys.* **2**, 275 (2006).
- [27] H. D. Rozenfeld, S. Havlin, and D. ben-Avraham, *New J. Phys.* **9**, 175 (2007).
- [28] H. D. Rozenfeld and D. ben-Avraham, *Phys. Rev. E* **75**, 061102 (2007).
- [29] A.-L. Barabási and R. Albert, *Science* **286**, 509 (1999).
- [30] C. Song, S. Havlin, H. A. Makse, *Nature* **433**, 392 (2005).
- [31] E. Domany, S. Alexander, D. Bensimon, and L. P. Kadanoff, *Phys. Rev. B* **28**, 3110 (1983).
- [32] M. G. Cosenza and R. Kapral, *Phys. Rev. A* **46**, 1850 (1992).
- [33] Z. Z. Zhang, J. H. Guan, W. L. Xie, Y. Qi, and S. G. Zhou, *EPL*, **86**, 10006 (2009).
- [34] Z. Z. Zhang, S. G. Zhou, L. C. Chen, and J. H. Guan, *Eur. Phys. J. B* **64**, 277 (2008).
- [35] Z. Z. Zhang, Y. Lin, and Y. J. Ma, *J. Phys. A* **44**, 075102 (2011).
- [36] A. Blumen, A. Jurjiu, Th. Koslowski, and Ch. von Ferber, *Phys. Rev. E* **67**, 061103 (2003).
- [37] A. Blumen, Ch. von Ferber, A. Jurjiu, and Th. Koslowski, *Macromolecules* **37**, 638 (2004).
- [38] Z. Z. Zhang, B. Wu, H. J. Zhang, S. G. Zhou, J. H. Guan, and Z. G. Wang, *Phys. Rev. E* **81**, 031118 (2010).
- [39] A. Jurjiu, A. Volta, and T. Beu, *Phys. Rev. E* **84**, 011801 (2011).
- [40] F. Chung, *Spectral Graph Theory* (American Mathematical Society, Providence, RI 1997).
- [41] F. Chung, L. Lu, and V. Vu, *Proc. Natl. Acad. Sci. U.S.A.* **100**, 6313 (2003).
- [42] H. Y. Chen and F. J. Zhang, *Discrete Appl. Math.* **155**, 654 (2007).
- [43] S. N. Dorogovtsev, A. V. Goltsev, J. F. F. Mendes, and A. N. Samukhin, *Phys. Rev. E* **68**, 046109 (2003).
- [44] D. Volchenkov and Ph. Blanchard, *Phys. Rev. E* **75**, 026104 (2007).
- [45] S. Hwang, C.-K. Yun, D.-S. Lee, B. Kahng, and D. Kim, *Phys. Rev. E* **82**, 056110 (2010).
- [46] C. Carretero-Campos, P. Bernaola-Galván, P. Ch. Ivanov, and P. Carpena, *Phys. Rev. E* **85**, 011139 (2012).
- [47] Z. Z. Zhang, H. X. Liu, B. Wu, and S. G. Zhou, *EPL* **90**, 68002 (2010).
- [48] Z. Z. Zhang, H. X. Liu, B. Wu, and T. Zou, *Phys. Rev. E* **83**, 016116 (2011).
- [49] Y. Lin, B. Wu, Z. Z. Zhang, and G. R. Chen, *J. Math. Phys.* **52**, 113303 (2011).
- [50] B. Wu, Z. Z. Zhang, and G. R. Chen, *J. Phys. A* **45**, 025102 (2012).
- [51] Z. Z. Zhang, B. Wu, and Y. Lin, *Physica A* **391**, 3342 (2012).
- [52] M. Moreau, G. Oshanin, O. Benichou, and M. Coppey, *Phys. Rev. E* **67**, 045104(R) (2003).
- [53] M. Moreau, G. Oshanin, O. Benichou, and M. Coppey, *Phys. Rev. E* **69**, 046101 (2004).
- [54] A. Bashan, R. P. Bartsch, Jan. W. Kantelhardt, S. Havlin, and P. Ch. Ivanov *Nat. Commun.* **3**, 702 (2012).